

Born–Infeld theory of electroweak and gravitational fields: Possible correction to Newton and Coulomb laws

Dmitriy Palatnik*

Abstract

In this note one suggests a possibility of direct observation of the θ -parameter, introduced in the Born–Infeld theory of electroweak and gravitational fields, developed in quant-ph/0202024. Namely, one may treat θ as a universal constant, responsible for correction to the Coulomb and Newton laws, allowing direct interaction between electrical charges and masses.

*SEL, Northwestern University Library, Evanston, IL 60208-3530 US; palatnik@ripco.com

1. In the note [3] a Born–Infeld (BI) [1] theory of electroweak and gravitational fields with the action invariant under θ -rotations of the Lorentz group, imposed on pairs (γ_a, π_a) , was developed. The following equations (slightly modified* from (6.5) and (6.6) in [3]) were obtained:

$$E' = \cosh \theta E + \frac{\hbar c}{\ell q} \sinh \theta Q; \quad (0.1)$$

$$Q' = \cosh \theta Q + \frac{\ell q}{\hbar c} \sinh \theta E. \quad (0.2)$$

Here

$$\ell^2 = \frac{10}{3\alpha}(1 + 2 \sin^2 \theta_W) L_p^2, \quad (0.3)$$

where L_p is the Planck's length, α is the fine structure constant for electron, and θ_W is the Weinberg mixing angle [2]. Equations (0.1) and (0.2) mean that if (E, Q) are possible solutions of field equations following from the theory for the energy and the electrical charge of, say, an electron, then (E', Q') are also possible solutions for the energy and the electrical charge of the same electron.

2. Consider, now, nonrelativistic case, so that $E \approx mc^2$, and $E' \approx m'c^2$. Suppose, that $\theta \ll 1$, so that one may neglect terms of order θ^2 . Substituting e and e' instead of Q and Q' respectively, one obtains from (0.1) and (0.2),

$$m' = m + \kappa^{-\frac{1}{2}} \theta e; \quad (0.4)$$

$$e' = e + \kappa^{\frac{1}{2}} \theta m. \quad (0.5)$$

Here

$$\kappa = \frac{q^2 \ell^2 c^2}{\hbar^2}. \quad (0.6)$$

One may interpret $\Delta m = \kappa^{-\frac{1}{2}} \theta e$ and $\Delta e = \kappa^{\frac{1}{2}} \theta m$ as contributions of electrical charge, e , and mass, m , to the observable mass, m' , and electrical charge, e' , respectively.

3. Keeping only leading terms of the action's expansion in ℓ [3], one would obtain standard equations for Coulomb's electrostatics and Newton's gravity in nonrelativistic case. Consider two electrical charges, (e, m) and (Q, M) . Then, observable quantities are (e', m') and (Q', M') . Here Q' and M' are expressed through Q and M similarly to (0.5) and (0.4). Then, one obtains for the Newton and Coulomb potential energy of two particles, separated at distance r ,

$$V(r) = \frac{e'Q' - km'M'}{r}. \quad (0.7)$$

* Here definition for the energy is $E = c \int \bar{\psi} p^0 \psi d\Sigma_0$, and for the electrical charge is $Q = q \int \bar{\psi} \gamma^0 \psi d\Sigma_0$, where ψ is a spinorial wave function of a particle, and q is electrical charge of electron (absolute value).

Here k is the Newton constant of gravitational interaction. Substituting ‘unprimed’ variables, one obtains for the potential energy,

$$V(r) = \frac{eQ - kmM + \sigma(eM + mQ)}{r}, \quad (0.8)$$

where

$$\sigma = \theta \left(\sqrt{\kappa} - \frac{k}{\sqrt{\kappa}} \right). \quad (0.9)$$

The new effect (proportional to σ) may be responsible for interaction between cosmic charged particles (protons, electrons, etc.) and the Earth’s ‘charge’, $Q_E = \sigma M_E$. Here M_E is the Earth’s mass. Assuming that $\sigma < 0$, one obtains that Earth’s mass contributes negative electric charge, which explains, why primary cosmic rays consist mainly from positively charged particles [4]. The same assumption explains the fairweather electric field (100 Volts per metre at surface of the Earth) [5]. Due to the same effect rotation of the Earth leads to electric currents, which may be responsible for Earth’s magnetic field.

REFERENCES

- [1] M. Born, L. Infeld, *Proc. Roy. Soc. A*, **144**, 425 (1934).
- [2] F. Halzen, A.D. Martin, *Quarks and Leptons*, N.Y. 1984.
- [3] D. Palatnik, LANL e-archive, quant-ph/0202024.
- [4] M. W. Friedlander, *Cosmic Rays*, Harvard U. Press, 1989.
- [5] L. Wahlin, *Atmospheric Electrostatics*, Research Studies Press, 1986.